arXiv:1604.07895v1 [math.CO] 27 Apr 2016

Fault-tolerance of balanced hypercubes with faulty vertices and faulty edges

Mei-Mei Gu
Department of Mathematics, Beijing Jiaotong University,
Beijing 100044, China
gum2012@bjtu.edu.cn

Rong-Xia Hao*
Department of Mathematics, Beijing Jiaotong University,
Beijing 100044, China
rxhao@bjtu.edu.cn

Abstract Let F_v (resp. F_e) be the set of faulty vertices (resp. faulty edges) in the n-dimensional balanced hypercube BH_n. Fault-tolerant Hamiltonian laceability in BH_n with at most $2n-2$ faulty edges is obtained in [Inform. Sci. 300 (2015) 20–27]. The existence of edge-Hamiltonian cycles in $BH_n - F_e$ for $|F_e| \leq 2n-2$ are gotten in [Appl. Math. Comput. 244 (2014) 447–456]. Up to now, almost all results about fault-tolerance in BH_n with only faulty vertices or only faulty edges. In this paper, we consider fault-tolerant cycle embedding of BH_n with both faulty vertices and faulty edges, and prove that there exists a fault-free cycle of length $2^{2n} - 2|F_v|$ in BH_n with $|F_v| + |F_e| \leq 2n-2$ and $|F_v| \leq n-1$ for $n \geq 2$. Since BH_n is a bipartite graph with two partite sets of equal size, the cycle of a length $2^{2n} - 2|F_v|$ is the longest in the worst-case.

Keywords Balanced hypercube; Cycle embedding; Fault tolerance; Interconnection network.

*This paper was accepted by Ars Combinatoria, March 1, 2016
*Corresponding author

1 Introduction

In a multiprocessor system, processors communicate by exchanging messages through an interconnection network whose topology often modeled by an undirected graph $G = (V, E)$, where every vertices in V corresponds to a processor, and every edge in E corresponds to a communication link.

The n-dimensional balanced hypercube BH_n, proposed by Wu and Huang [16], is an important class of generalizations of the popular hypercube interconnection network for parallel computing. It has many desirable properties, such as bipartite, regularity, recursive structure, vertex-transitive [10] and edge transitive [25], as the hypercube. However, the balanced hypercube is superior to the hypercube in a sense that it has a smaller diameter than that of the hypercube and supports an efficient reconfiguration without changing the adjacent relationship among tasks [16]. More desired properties of balanced hypercubes have been shown in the literature. Let BH_n be an n-dimensional balanced hypercube. Wu and Huang [10] proved that BH_n is bipartite graph and vertex transitive. Huang and Wu [11] studied resource placement problem in BH_n. Zhou et al. [25] obtained that BH_n is edge transitive and determined the cyclic connectivity and cyclic edge-connectivity of BH_n. Xu et al. [18] showed that BH_n is edge-pancyclic and Hamiltonian laceable. Yang [19] further showed that BH_n is bipanconnected for all $n \geq 1$. Yang [20] determined super connectivity and super edge connectivity of BH_n. Yang [21] also gave the conditional diagnosability of BH_n under the PMC model. Lü et al. [13] obtained (conditional) matching preclusion number of BH_n. Lü and Zhang [14] further proved that BH_n is hyper-Hamiltonian laceable. Cheng et al. [5] showed that there exist two vertex-disjoint path cover in BH_n.

Since faulty vertices and faulty edges may happen in the real world networks, the fault-tolerant of networks has been proposed, see [7, 9, 15, 17, 24] etc.. There are some results about the fault-tolerance of BH_n. Let F_v (resp. F_e) be the set of faulty vertices (resp. faulty edges) in the n-dimensional balanced hypercube BH_n. As examples, Zhou et al. [26] derived that BH_n is $(2n-2)$-edge-fault Hamiltonian laceable for $n \geq 2$, i.e., if at most $2n-2$ faulty edges occurs, the resulting graph remains Hamiltonian laceable. Cheng et al. in [4] proved that if $|F_v| \leq n-1$ faulty vertices occurs, every fault-free edge of $BH_n - F_v$ lies on a fault-free cycle of every even length from 4 to $2^{2n} - 2|F_v|$. Cheng et al. in [3] derived that BH_n is $(2n-3)$-edge-fault-bipancyclic, that is, there are at most $|F_e| \leq 2n-3$ faulty edges, every fault-free edge of $BH_n - F_e$ lies on a fault-free cycle of every even length from 4 to 2^{2n}. Hao et al. [8] gave the existence of $(2n-2)$- edge-fault-tolerant cycle embedding in BH_n. But up to now, almost all results about fault-tolerance in BH_n with only faulty vertices or only faulty edges.

In this paper, we consider cycle embedding of BH_n with both faulty vertices and faulty edges, and prove that there exists a fault-free cycle of length $2^{2n} - 2|F_v|$ in BH_n with $|F_v| + |F_e| \leq 2n - 2$ and $|F_v| \leq n - 1$ for $n \geq 2$. Since BH_n is a bipartite graph with two partite sets of equal size, the cycle of length $2^{2n} - 2|F_v|$ is the longest in the worst-case. This paper improves the results about the longest fault-free cycle in BH_n with only faulty vertices or only faulty edges.

The rest of this paper is organized as follows. Section 2 presents some necessary definitions of graphs and properties of balanced hypercubes as preliminaries. The proof of our main result is given in Section 3. Section 4 concludes this paper.

2 Preliminaries

Let $G = (V, E)$ be a simple undirected graph, where V is the vertex-set of G and E is the edge set of G. The *neighborhood* of a vertex u in G is the set of all vertices adjacent to u in $V(G)$, denoted by $N_G(u)$. The cardinality $|N_G(u)|$ represents the *degree* of u in G, denoted by $d_G(u)$. A *path* P, denoted by $P = \langle u_0, u_1, \ldots, u_k \rangle$, is a sequence of vertices where two successive vertices are adjacent in G. u_0 and u_k are called the *end-vertices* of a path P. A $u - v$ *path* which is denoted by $P[u, v]$ is a path with endpoints u and v. For $0 \leq i < j \leq k$, let $P = \langle u_0, u_1, \ldots, u_k \rangle = \langle u_0, u_1, \ldots, u_i, P[u_i, u_j], u_j, \ldots, u_k \rangle$, where $P[u_i, u_j] = \langle u_i, u_{i+1}, \ldots, u_j \rangle$. A *cycle* is a path that two ends are the same vertex. A cycle (resp. path) is a *Hamiltonian cycle* (resp. *Hamiltonian path*) if it traverses all the vertices of G exactly once. The *length* of a path is the number of edges in it.

A graph G is said to be *Hamiltonian connected* if there exists a Hamiltonian path between any two vertices of G. It is easy to see that any bipartite graph with at least three vertices is not Hamiltonian connected. A bipartite graph G is *Hamiltonian laceable* (resp. *strongly Hamiltonian laceable*) if there exists a of length $|V(G)| - 1$ (resp. $|V(G)| - 2$) between two arbitrary vertices from different (resp. same) partite sets. A bipartite graph G with bipartition $V_0 \cup V_1$ is *hyper-Hamiltonian laceable* if it is Hamiltonian laceable and, for any vertex $v \in V_i$ ($i \in \{0, 1\}$), there is a Hamiltonian path in $G - v$ between any pair of vertices in V_{1-i}.

An *n-dimensional balanced hypercube*, denoted by BH_n, is proposed by Wu and Huang [16] which is defined as an undirected simple graph.

Definition 2.1 *An n-dimensional balanced hypercube BH_n with $n \geq 1$ has 2^{2n} vertices with addresses $(a_0, a_1, \ldots, a_{n-1})$, where $a_i \in \{0, 1, 2, 3\}$ for each $0 \leq i \leq n - 1$. Each vertex $(a_0, a_1, \ldots, a_{n-1})$ is adjacent to the following $2n$ vertices:*

$$\begin{cases} ((a_0 \pm 1)(\text{mod } 4), a_1, \ldots, a_{n-1}); \\ ((a_0 \pm 1)(\text{mod } 4), a_1, \ldots, a_{j-1}, (a_j + (-1)^{a_0})(\text{mod } 4), a_{j+1}, \ldots, a_{n-1}), \end{cases}$$

where j is an integer with $1 \leq j \leq n-1$.

'+' and '$-$' are the operation with "(mod 4)". For notation convenience, "(mod 4)" is omitted. The balanced hypercube BH_n can also be recursively defined.

Definition 2.2 *BH_n can be recursively constructed as follows:*

(1) *BH_1 is a cycle consisting of 4 vertices labeled as $0, 1, 2, 3$ respectively.*

(2) *For $n \geq 2$, BH_n consists of four copies of BH_{n-1}, denoted by BH_{n-1}^i for each $i \in \{0, 1, 2, 3\}$. Each vertex $(a_0, a_1, \ldots, a_{n-2}, i)$ in BH_{n-1}^i has two extra adjacent vertices which are also called extra neighbors:*

(2.1) *$(a_0 \pm 1, a_1, \ldots, a_{n-2}, i+1)$ in BH_{n-1}^{i+1} if a_0 is even.*

(2.2) *$(a_0 \pm 1, a_1, \ldots, a_{n-2}, i-1)$ in BH_{n-1}^{i-1} if a_0 is odd.*

BH_1 and BH_2 are illustrated in Figure 1.



Figure 1: BH_1 and BH_2

The first element a_0 of vertex $(a_0, a_1, \ldots, a_{n-2}, a_{n-1})$ is called the *inner index*, and the other elements a_i, for all $1 \leq i \leq n-1$, are called *i-dimensional index*. By [23, Lemma 5], for each vertex u of BH_n, there exists a unique vertex, denoted by u^*, such that $N_{BH_n}(u) = N_{BH_n}(u^*)$.

Since BH_n is a bipartite graph, we refer to a vertex with an odd inner index as a *black vertex* and a vertex with an even inner index as a *white vertex*. If two adjacent vertices u and v differ in only the inner index, the edge $e = (u, v)$ is said to be 0-*dimensional* and v is a

0-*dimensional neighbor* of u. Let $[n] = \{0, 1, \ldots, n-1\}$ (but not general meaning $\{1, 2, \ldots, n\}$). Likewise if two adjacent vertices u and v not only differ in the inner index, but also differ in some j-dimensional index ($j \in [n]$), the edge $e = (u, v)$ is said to be *j-dimensional*, and v is the *j-dimensional neighbor* of u. Let E_j, where $j \in [n]$, denote the set of all j-dimensional edges. For $i \in \{0, 1, 2, 3\}$ and $j \in [n]$, we use $BH_{n-1}^{j,i}$ to denote $(n-1)$-dimensional sub-balanced hypercubes of the BH_n induced by all vertices labeled by $(a_0, a_1, \ldots, a_{j-1}, i, a_{j+1}, \ldots, a_{n-2}, a_{n-1})$. Obviously, $BH_n - E_j = \bigcup_{i=0}^{3} BH_{n-1}^{j,i}$ and $BH_{n-1}^{j,i} \cong BH_{n-1}$. The edges between $BH_{n-1}^{j,i}$ and $BH_{n-1}^{j,i+1}$ are called *j-dimensional crossing edges*, where $i \in \{0, 1, 2, 3\}$ and $j \in \{1, 2, \ldots, n-1\}$. If $j = n-1$, $BH_{n-1}^{j,i}$ and E_j are denoted by BH_{n-1}^{i} and E_c, respectively, where $i \in \{0, 1, 2, 3\}$ and $(n-1)$-dimension crossing edges are called *crossing edges*. Let E_0 be the set of all 0-dimensional edges, Lü et al. [13] obtained that $BH_n - E_0$ has four components, each component is isomorphic to BH_{n-1}.

Some known basic properties of BH_n are given as follows.

Lemma 2.3 ([16]) *The balanced hypercube BH_n is bipartite and vertex transitive.*

Lemma 2.4 *(*[25]) *The balanced hypercube BH_n is edge transitive.*

Lemma 2.5 ([18]) *The balanced hypercube BH_n is edge-bipancyclic and Hamiltonian laceable for all $n \geq 1$.*

Lemma 2.6 *(*[19]) *The balanced hypercube BH_n is bipanconnected for all $n \geq 1$.*

Lemma 2.7 ([14]) *The balanced hypercube BH_n is hyper-Hamiltonian laceable for $n \geq 1$. Thus, BH_n is strongly Hamiltonian laceability.*

Lemma 2.8 ([5]) *Let X and Y be two distinct partite sets of BH_n. Assume u and x are two different vertices in X, v and y are two different vertices in Y. Then there exist two vertex-disjoint paths $P[x, y]$ and $R[u, v]$, and $V(P[x, y]) \cup V(R[u, v]) = V(BH_n)$.*

Lemma 2.9 (1) ([18]) *Let $e = (u, v)$ be an edge of BH_n. e is contained in a cycle C of length 8 in BH_n such that $|E(C) \cap E(BH_{n-1}^{i})| = 1$, where $i = 0, 1, 2, 3$.*

(2) ([4]) *Let $e = (u, v)$ be an edge of BH_n. If e is along dimension j ($1 \leq j \leq n-1$), then there are $2n-2$ internal vertex-disjoint paths of length 7 joining u and v such that each path has only one edge in each $BH_{n-1}^{j,i}$, where $i \in \{0, 1, 2, 3\}$.*

(3) ([4]) *Let $e = (x, y)$ be any edge in $BH^{j,0}_{n-1}$, then there exist two internal vertex-disjoint paths $\langle x, y_1, x_1, y_2, x_2, y_3, x_3, y\rangle$ and $\langle x, y'_1, x'_1, y'_2, x'_2, y'_3, x'_3, y\rangle$ in BH_n such that $(x_i, y_i), (x'_i, y'_i) \in E(BH^{j,i}_{n-1})$, where $1 \leq j \leq n-1$ and $i = 1, 2, 3$.*

Lemma 2.10 ([8]) *Let $A = \{e_1, e_2, \ldots, e_k\}$ for $k \geq 2$ be the set of j-dimensional edges in BH^i_{n-1}, where $j \in [n]$ and $i \in \{0, 1, 2, 3\}$. Then there are cycles $D_1, D_2, \ldots, D_k$ which contain $e_1, e_2, \ldots, e_k$, respectively, such that the length of each cycle is 8, $D_1, D_2, \ldots, D_k$ are edge disjoint and $|E(D_m) \cap E(BH^t_{n-1})| = 1$ for every $m \in \{1, 2, \ldots, k\}$ and $t \in \{0, 1, 2, 3\}$. Moreover, for any $i, j \in [k]$, if $e_i \cap e_j = \emptyset$, then D_i, D_j are also vertex-disjoint. If $e_i \cap e_j = \{u\}$, then D_i, D_j are vertex-disjoint except a vertex u.*

For the fault tolerance of BH_n, Cheng et al. [3] and [4] derived the following two results, respectively..

Lemma 2.11 ([3]) *Let F_e be the set of faulty vertices in BH_n with $|F_e| \leq 2n-3$. Then every fault-free edge of $BH_n - F_e$ lies on a fault-free cycle of every even length from 4 to 2^{2n} inclusive, where $n \geq 2$.*

Lemma 2.12 ([4]) *Let F_v be the set of faulty vertices in BH_n with $|F_v| \leq n-1$. Then every fault-free edge of $BH_n - F_v$ lies on a fault-free cycle of every even length from 4 to $2^{2n} - 2|F_v|$ inclusive, where $n \geq 1$.*

Hao et al. [8] and Zhou et al. [26] obtained the following two results, respectively.

Lemma 2.13 ([8]) *Let F_e be a faulty edge set of the balanced hypercube BH_n with $|F_e| \leq 2n-2$, $n \geq 1$, then there exists a fault-free Hamiltonian path between any two adjacent vertices x and y in BH_n.*

Lemma 2.14 ([26]) *BH_n is $(2n-2)$-edge-fault Hamiltonian laceable for $n \geq 2$.*

3 Main results

In this section, we consider cycle embedding in faulty balanced hypercube BH_n. For this purpose, we need the following lemma.

Lemma 3.1 *Let F_v (resp. F_e) be the set of faulty vertices (resp. faulty edges) with $|F_v| + |F_e| \leq n-1$ in BH_n for $n \geq 1$. Then, there is a fault-free path in BH_n whose length is $2^{2n} - 2|F_v| - 1$ between any two adjacent vertices x and y in BH_n.*

Proof. Assume x is a white vertex and y is a black vertex and $(x, y) \in E(BH_n)$. Recall that $f_e = |F_e|$, $f_v = |F_v|$, $f = f_e + f_v$. We only need to prove the result holds for $f = n - 1$. We show the lemma by induction on n. If $n = 1$, $f = 0$, BH_1 is a cycle of length 4, so the result is right. Now we consider $n = 2$. If $f_e = 0$, then $f_v = 1$. Since x and y are adjacent, regard (x, y) as a fault-free edge, by Lemma 2.12, the edge (x, y) lies in a fault-free cycle of length $2^4 - 2$, this implies there exists a fault-free path $P[x, y]$ of length $2^4 - 2 - 1 = 13$. If $f_v = 0$, then $f_e = 1$. Since x and y are adjacent, by Lemma 2.13, there exists a fault-free Hamiltonian $x - y$ path of length $2^4 - 1 = 15$. The lemma holds for $n = 2$.

Assume that the lemma holds for BH_{n-1} for $n \geq 3$. Now we consider BH_n. If BH_n has no faulty edges, then $f_v \leq n - 1$. By Lemma 2.12, there exists a fault-free path of length $2^{2n} - 2f_v - 1$ between x and y. So we assume that there is at least one faulty edge. Without loss of generality, suppose there exists an $(n-1)$-dimensional faulty edge. We can divide BH_n into BH^i_{n-1}, $i = 0, 1, 2, 3$, along $(n - 1)$-dimension. For $i \in \{0, 1, 2, 3\}$, let $f^i_e = |F_e \cap E(BH^i_{n-1})|$ and $f^i_v = |F_v \cap V(BH^i_{n-1})|$. Moreover, we have that $0 \leq f^i_e + f^i_v \leq n - 2$ for each $i \in \{0, 1, 2, 3\}$. There are the following two cases.

Case 1. x and y are in the same BH^i_{n-1} for some $i \in \{0, 1, 2, 3\}$. Without loss of generality, let $i = 0$.

By the inductive hypothesis, there exists a fault-free $x - y$ path P_0 in BH^0_{n-1} of length $2^{2(n-1)} - 2f^0_v - 1$. We claim that there exists at least one $k \in \{0, 1, \ldots, n - 2\}$ such that P_0 contains at least $2n - 1$ k-dimensional edges. Otherwise, for every $k \in \{0, 1, \ldots, n-2\}$, P_0 contains at most $2n-2$ k-dimensional edges. As a result, P_0 contains at most $(n-1)(2n-2)$ edges. Since $|P_0| = 2^{2(n-1)} - 2f^0_v - 1 \geq 2^{2(n-1)} - 2n + 1 > (n - 1)(2n - 2)$ for $n \geq 3$, it is a contradiction. Since $f = n - 1 \leq 2n - 1$ for $n \geq 3$, by Lemma 2.10, we can find a k-dimensional edge (u_0, v_0) (assume u_0 is white, v_0 is black) on P_0, such that (u_0, v_0) is contained in a fault-free 8-cycle C, say $C = \langle u_0, v_0, u_3, v_3, u_2, v_2, u_1, v_1, u_0\rangle$, where $(u_i, v_i) \in E(BH^i_{n-1})$ for $i \in \{0, 1, 2, 3\}$.

By the inductive hypothesis in BH^i_{n-1}, for each $i \in \{1, 2, 3\}$, there exists a fault-free $u_i - v_i$ path P_i in BH^i_{n-1} of length $2^{2(n-1)} - 2f^i_v - 1$. Therefore, $\langle x, P[x, u_0], u_0, v_1, P_1, u_1, v_2, P_2,$
$u_2, v_3, P_3, u_3, v_0, P[v_0, y], y\rangle$ is a desired fault-free $x - y$ path in BH_n of length $\Sigma^3_{i=0}(2^{2(n-1)} - 2f^i_v - 1) + 3 = 2^{2n} - 2f_v - 1$.

Case 2. x and y are in two distinct BH^i_{n-1}'s, where $i \in \{0, 1, 2, 3\} = [4]$.

Without loss of generality, suppose $x \in V(BH^0_{n-1})$. Since x and y are adjacent, $y \in V(BH^1_{n-1})$. By Lemma 2.9(2) and $f \leq n-1 \leq 2n-2$ for $n \geq 3$, there exists a fault-free $x-y$ path P, say $P = \langle x, v_0, u_3, v_3, u_2, v_2, u_1, y\rangle$, of length 7 such that P has only one edge in BH^i_{n-1} for each $i = 0, 1, 2, 3$.

By the inductive hypothesis, there exists a fault-free $x - v_0$ (resp. $u_3 - v_3$, $u_2 - v_2$ and $u_1 - y$) path P_0 (resp. P_3, P_2 and P_1) in BH^0_{n-1} (resp. BH^3_{n-1}, BH^2_{n-1} and BH^1_{n-1}) of length $2^{2(n-1)} - 2f^0_v - 1$ (resp. $2^{2(n-1)} - 2f^3_v - 1$, $2^{2(n-1)} - 2f^2_v - 1$ and $2^{2(n-1)} - 2f^1_v - 1$). A fault-free $x - y$ path of BH_n can be constructed as $\langle x, P_0, v_0, u_3, P_3, v_3, u_2, P_2, v_2, u_1, P_1, y\rangle$ of length $\Sigma^3_{i=0}(2^{2(n-1)} - 2f^i_v - 1) + 3 = 2^{2n} - 2f_v - 1$.

By the above cases, the proof is complete. □

Lemma 3.2 *Let v be a faulty vertex and e be a faulty edge in BH_2. Then there exists a fault-free cycle of length 14 in BH_2.*

Proof. By Lemma 2.4, BH_2 is edge transitive, we may assume that $a = ((0,0),(1,1))$ is a faulty edge. For each possible faulty vertex $v \in V(BH_2)$, we can find a desired cycle. All the cases are listed in Table 1. □

$v = (0,0)$ or $(1,0)$
$\langle(3,0),(2,0),(3,1),(0,1),(1,1),(2,1),(1,2),(2,2),(3,2),(0,2),(1,3),(2,3),(3,3),(0,3),(3,0)\rangle$
$v = (3,0)$ or $(2,0)$
$\langle(0,0),(1,0),(0,3),(3,3),(2,3),(1,3),(0,2),(3,2),(2,2),(1,2),(2,1),(1,1),(0,1),(3,1),(0,0)\rangle$
$v = (0,1)$ or $(1,1)$
$\langle(0,0),(1,0),(2,0),(3,1),(2,1),(1,2),(2,2),(3,2),(0,2),(1,3),(2,3),(3,3),(0,3),(3,0),(0,0)\rangle$
$v = (3,1)$ or $(2,1)$
$\langle(0,0),(1,0),(2,0),(1,1),(0,1),(3,2),(2,2),(1,2),(0,2),(1,3),(2,3),(3,3),(0,3),(3,0),(0,0)\rangle$
$v = (0,3)$ or $(1,3)$
$\langle(0,0),(3,0),(2,0),(1,0),(2,3),(3,3),(2,2),(3,2),(0,2),(1,2),(2,1),(1,1),(0,1),(3,1),(0,0)\rangle$
$v = (3,3)$ or $(2,3)$
$\langle(0,0),(3,0),(2,0),(1,0),(0,3),(1,3),(0,2),(3,2),(2,2),(1,2),(2,1),(1,1),(0,1),(3,1),(0,0)\rangle$
$v = (0,2)$ or $(1,2)$
$\langle(0,0),(1,0),(2,0),(3,0),(0,3),(3,3),(2,3),(1,3),(2,2),(3,2),(0,1),(1,1),(2,1),(3,1),(0,0)\rangle$
$v = (3,2)$ or $(2,2)$
$\langle(0,0),(1,0),(2,0),(3,0),(0,3),(3,3),(2,3),(1,3),(0,2),(1,2),(2,1),(1,1),(0,1),(3,1),(0,0)\rangle$

Table 1: Desired cycles of length 14 in BH_2 with a faulty vertex v and a faulty edge ((0,0),(1,1))

The following is our main result.

Theorem 3.3 *Let F_v and F_e be the set of faulty vertices and faulty edges, respectively, with $|F_v| \leq n-1$ and $|F_v| + |F_e| \leq 2n-2$ in BH_n, where $n \geq 2$. Then there exists a fault-free cycle of length $2^{2n} - 2|F_v|$ in BH_n.*

Proof. We prove this theorem by induction on n. Recall that E_j is the set of j-dimensional edges in BH_n for $j \in [n]$. Since $|F_v| + |F_e| \leq 2n-2$, there exists a $j \in [n]$ such that $|F_e \cap E_j| \leq 1$. Without loss of generality, let $j = n-1$. Then we can divide BH_n into four components, say BH^i_{n-1}, $i = 0, 1, 2, 3$, along $(n-1)$-dimension. Let $F_c = F_e \cap E_{n-1}$, then $f_c = |F_c| \leq 1$. Recall that $f_v = |F_v|$, $f_e = |F_e|$, $f = f_v + f_e$, $f^i_e = |F_e \cap E(BH^i_{n-1})|$ and $f^i_v = |F_v \cap V(BH^i_{n-1})|$, so $f_i = f^i_e + f^i_v$ for $i \in \{0, 1, 2, 3\}$.

If $n = 2$, then $f_v + f_e \leq 2$ and $f_v \leq 1$. If $f_v = 1$ and $f_e = 0$, then by Lemma 2.12, the result is right. If $f_v = 1$ and $f_e = 1$, the result follows from Lemma 3.2. If $f_v = 0$ and $f_e \leq 2$. Choose a fault-free edge (u, v), by Lemma 2.13, there exists a fault-free Hamiltonian path P between u and v in BH_2, then $\langle u, P, v, u\rangle$ is the desired cycle of BH_2. The theorem is true for $n = 2$.

For $n \geq 3$, we assume that the theorem is true in BH_m with $f_v + f_e \leq 2m-2$ and $f_v \leq m-1$ for every integer $2 \leq m < n$. Now we consider BH_n as follows. If BH_n has no faulty vertices, then $f_v = 0$ and $f_e \leq 2n-2$. Choose a fault-free edge (x, y), since $f_e \leq 2n-2$, by Lemma 2.13, there exists a fault-free Hamiltonian path P from x to y in BH_n. Let $C = \langle x, P, y, x\rangle$. Clearly, $|E(C)| = 2^{2n} = 2^{2n} - 2f_v$. So we assume $1 \leq f_v \leq n-1$ in the following. We only need to consider the following three cases.

Case 1. For each $i \in \{0, 1, 2, 3\}$, $0 \leq f_i \leq 2n-4$.

Subcase 1.1. For each $i \in \{0, 1, 2, 3\}$, $0 \leq f_i \leq n-2$.

By the inductive hypothesis, there exists a cycle C_0 in BH^0_{n-1} of length $2^{2(n-1)} - 2f^0_v$. Select an $(u_0, v_0) \in E(C_0)$ and assume u_0 is white. By Lemma 2.9(3), since $f_c \leq 1$, (u_0, v_0) is contained in two 8-cycles which has only one common edge (u_0, v_0), so we can find a 8-cycle C, say $C = \langle u_0, v_0, u_3, v_3, u_2, v_2, u_1, v_1, u_0\rangle$, where $(u_i, v_i) \in E(BH^i_{n-1})$ for $i \in \{0, 1, 2, 3\}$, such that the crossing edge (u_i, v_{i+1}) for each $i \in \{0, 1, 2, 3\}$ is fault-free, where $v_4 = v_0$. Since $f_i \leq n-2$ and $BH^i_{n-1} \cong BH_{n-1}$ for $i \in \{1, 2, 3\}$, by Lemma 3.1, there exists a fault-free $u_i - v_i$ path P_i of length $2^{2(n-1)} - 2f^i_v - 1$ in BH^i_{n-1}. Let $P_0 = C_0 - (u_0, v_0)$. So $\langle u_0, v_1, P_1, u_1, v_2, P_2, u_2, v_3, P_3, u_3, v_0, P_0, u_0\rangle$ is a desired fault-free cycle in BH_n of length $2^{2n} - 2f_v$.

Subcase 1.2. There exists only one $i \in \{0, 1, 2, 3\}$ such that $n-1 \leq f_i \leq 2n-4$. Without loss of generality, let $i = 0$. It implies that $0 \leq f_j \leq n-2$ for all $j \in \{1, 2, 3\}$.

Subcase 1.2.1. $n-1 \leq f_0 \leq 2n-4$, $f^0_v \leq n-2$.

By the inductive hypothesis, there exists a fault-free cycle C_0 in BH^0_{n-1} of length $2^{2(n-1)} - 2f^0_v$. By the similar discussions as Case 1 in the proof of Lemma 3.1, we can find an edge (u_0, v_0) (assume u_0 is white, v_0 is black) on C_0, such that (u_0, v_0) is contained in a fault-free 8-cycle C, say $C = \langle u_0, v_0, u_3, v_3, u_2, v_2, u_1, v_1, u_0\rangle$, where $(u_i, v_i) \in E(BH^i_{n-1})$ for $i \in \{1, 2, 3\}$. Since $0 \leq f_i \leq n-2$ for $i \in \{1, 2, 3\}$, by Lemma 3.1 in BH^i_{n-1}, there

exists a fault-free $u_i - v_i$ path P_i in BH^i_{n-1} of length $2^{2(n-1)} - 2f^i_v - 1$. Let $P_0 = C_0 - (u_0, v_0)$. Therefore, $\langle u_0, v_1, P_1, u_1, v_2, P_2, u_2, v_3, P_3, u_3, v_0, P_0, u_0\rangle$ is a desired fault-free cycle in BH_n of length $2^{2n} - 2f_v$.

Subcase 1.2.2. $n-1 \leq f_0 \leq 2n-4$, $f^0_v = n-1$.

It implies that $f^j_v = 0$ and $f^j_e \leq f - f_v \leq n-1$ for each $j \in \{1,2,3\}$. We regard one faulty vertex in BH^0_{n-1} as a fault-free vertex, so $f'_v = f^0_v - 1 = f_v - 1 = n-2$. By the inductive hypothesis, there exists a cycle C_0 in BH^0_{n-1} of length $2^{2(n-1)} - 2f'_v$, where $f'_v = f^0_v - 1$. Note that C_0 contains at most one faulty vertex.

If C_0 does not contain any faulty vertex, choose an edge, say $(u_0, v_0) \in E(C_0)$ (assume u_0 is white, v_0 is black). By Lemma 2.9(3) and $f_c \leq 1$, (u_0, v_0) is contained in a cycle C, say $C = \langle v_0, u_0, v_1, u_1, v_2, u_2, v_3, u_3, v_0\rangle$, of length 8 in BH_n such that $E(C) \cap E(BH^i_{n-1}) = \{(u_i, v_i)\}$ and (u_i, v_{i+1}) is fault-free for each $i \in \{0,1,2,3\}$. Since for any $i \in \{1,2\}$, $f^i_v = 0$ and $f^i_e \leq n-1 \leq 2n-2$ for $n \geq 3$, by Lemma 2.14, there exists a fault-free path P_i between u_i and v_i in BH^i_{n-1} whose length is $2^{2(n-1)} - 1$. Since $f^3_v = 0$ and $f^3_e \leq n-1 \leq 2n-3$ for $n \geq 3$, by Lemma 2.11, there exists a fault-free cycle C_3 of length $2^{2(n-1)} - 2$ passing through (u_3, v_3). Let $P_i = C_i - (u_i, v_i)$ for $i = 0, 3$. A desired fault-free cycle of BH_n can be constructed as $\langle u_0, v_1, P_1, u_1, v_2, P_2, u_2, v_3, P_3, u_3, v_0, P_0, u_0\rangle$ whose length is $2^{2(n-1)} - 2(f^0_v - 1) - 1 + 2(2^{2(n-1)} - 1) + (2^{2(n-1)} - 3) + 4 = 2^{2n} - 2f^0_v = 2^{2n} - 2f_v$.

Now assume that C_0 contains one faulty vertex u_0 and u_0 is black. (If C_0 contains a white faulty vertex, the discussions are similar, so it is omitted.) Let a_0 and b_0 be two vertices adjacent to u_0 on C_0. Let $c_0 \in C_0$ be a neighbor of b_0. Recall that each vertex has two extra neighbors and $f_c \leq 1$, we can choose the extra neighbors $c_1 \in V(BH^1_{n-1})$ and $a_3 \in V(BH^3_{n-1})$ of a_0 and c_0, respectively, such that both (a_0, c_1) and (c_0, a_3) are fault-free. For $i \in \{1,2\}$, let $(a_i, c_{i+1}) \in E(BH_n)$ be a fault-free edge, where $a_i \in V(BH^i_{n-1})$ be a white vertex and $c_{i+1} \in V(BH^{i+1}_{n-1})$ be a black vertex. For $i \in \{1,2,3\}$, $f^i_e \leq n-1 \leq 2n-4$ for $n \geq 3$ and $f^i_v = 0$, by Lemma 2.14, there exists a fault-free path P_i of length $2^{2(n-1)} - 1$ between a_i and c_i in BH^i_{n-1}. Let $P_0 = C_0 - \{u_0, b_0\}$. A desired fault-free cycle of BH_n can be constructed as $\langle a_0, P_0, c_0, a_3, P_3, c_3, a_2, P_2, c_2, a_1, P_1, c_1, a_0\rangle$ of length $2^{2n} - 2f_v$ (see Figure 2).

Subcase 1.3. There exist two distinct $i, k \in \{0,1,2,3\}$ such that $n-1 \leq f_i \leq 2n-4$ and $n-1 \leq f_k \leq 2n-4$. Without loss of generality, assume that $i = 0$. Since $f \leq 2n-2$, it implies that $f_0 = f_k = n-1$, $f_c = |F_c| = 0$ and $f_t = 0$ for any $t \in \{1,2,3\} - \{k\}$.

Subcase 1.3.1. $f^0_v \leq n-2$ and $f^k_v \leq n-2$.

By the inductive hypothesis, there exists a cycle C_0 (resp. C_k) in BH^0_{n-1} (resp. BH^k_{n-1}) of length $2^{2(n-1)} - 2f^0_v$ (resp. $2^{2(n-1)} - 2f^k_v$).



Figure 2: Illustration of 1.2.2



Figure 3: Illustration of 1.3.1.1

Subcase 1.3.1.1. $k = 1$ or $k = 3$.

Since the discussion for the two situations are similar, we may assume $k = 1$.

Note that C_0 has $2^{2(n-1)} - 2f_v^0$ vertices, half is black and half is white, so there are $2^{2n-3} - f_v^0$ white vertices. We claim that there exist at least one white vertex in C_0 such that its extra neighbor in C_1. Since for each white vertex, say u, in C_0, there exists at most one other vertex u^* in C_0, which has the same extra neighbors as u. So vertices in C_0 have at least $(2^{2n-3} - f_v^0)/2$ extra neighbors in C_1. By $(2^{2n-3} - f_v^0)/2 > n - 1 = f_1$ because of $f_v^0 \leq n - 2$ and $n \geq 3$, there exists a white vertex $u_0 \in V(C_0)$ such that u_0 has an extra neighbor, say v_1, on C_1. See Figure 3.

Let v_0 be a neighbor of u_0 in C_0 and u_1 be a neighbor of v_1 in C_1. Let $u_3 \in V(BH^3_{n-1})$ (resp. $v_2 \in V(BH^2_{n-1})$) be an extra neighbor of v_0 (resp. u_1). Let $(v_2, u_3) \in E(BH_n)$, where $v_2 \in V(BH^2_{n-1})$ be a black vertex and $u_3 \in V(BH^3_{n-1})$ be a white vertex. Since $f_t = 0$ for $t \in \{2, 3\}$, by Lemma 2.5, there exists a fault-free path P_t of length $2^{2(n-1)} - 1$ between u_t and v_t in BH^t_{n-1}. Let $P_i = C_i - (u_i, v_i)$ for $i = 0, 1$, then $\langle u_0, P_0, v_0, u_3, P_3, v_3, u_2, P_2, v_2, u_1, P_1, v_1, u_0\rangle$ is a fault-free cycle of length $2^{2n} - 2f_v$ in BH_n.

Subcase 1.3.1.2. $k = 2$.

Let $(u_0, v_0) \in C_0$ and assume u_0 is white. Choose an edge $(u_2, v_2) \in E(C_2)$, assume u_2 is white. Let $v_1 \in V(BH^1_{n-1})$ (resp. $u_1 \in V(BH^1_{n-1})$, $v_3 \in V(BH^3_{n-1})$ and $u_3 \in V(BH^3_{n-1})$) be an extra neighbor of u_0 (resp. v_2, u_2 and v_0). Since $f_t = 0$ for $t \in \{1, 3\}$, by Lemma 2.5, there exists a fault-free path P_t of length $2^{2(n-1)} - 1$ between u_t and v_t in BH^t_{n-1}. Let $P_i = C_i - (u_i, v_i)$ for $i = 0, 2$, then $\langle u_0, P_0, v_0, u_3, P_3, v_3, u_2, P_2, v_2, u_1, P_1, v_1, u_0\rangle$ is a desired fault-free cycle of length $2^{2n} - 2f_v$ in BH_n.

Subcase 1.3.2. $f_v^0 = n-1$. (The discussion is similar for $f_v^k = n-1$.)

It implies $f_v^k = 0$ and $f_e^k = n-1$. We regard one faulty vertex in BH_{n-1}^0 as fault-free, $f_v' = f_v^0 - 1 = n-2$. By the inductive hypothesis, there exists a cycle C_0 in BH_{n-1}^0 of length $2^{2(n-1)} - 2f_v'$, where $f_v' = f_v^0 - 1 = f_v - 1$. Note that C_0 contains at most one faulty vertex. Without loss of generality, assume that C_0 contains one faulty vertex u_0 and u_0 is black. Let a_0 and b_0 be two vertices adjacent to u_0 in C_0. Let $c_0 \in C_0$ be a neighbor of b_0. The desired cycle of BH_n can be constructed by the similar discussions as Subcase 1.2.2, the details are omitted.

Case 2. There exists one $i \in \{0,1,2,3\}$ such that $f_i = 2n-3$.

Without loss of generality, let $i = 0$, i.e., $f_0 = 2n-3$. It implies that $f_t \leq f - f_0 \leq 1$ for all $t \in \{1,2,3\}$. We consider the following two subcases.

Subcase 2.1. $f_v^0 \leq n-2$.

Recall that $f_e^0 = f_0 - f_v^0 \geq 2n-3-(n-2) = n-1 \geq 2$ for $n \geq 3$. Choose any faulty edge, say $(u_0, v_0) \in E(BH_{n-1}^0)$, regard (u_0, v_0) as a fault-free edge temporarily, then $f_0 - 1 \leq 2n-4$. By the inductive hypothesis, there exists a cycle C_0 of length $2^{2(n-1)} - 2f_v^0$ in BH_{n-1}^0 such that C_0 contains at most one faulty edge (u_0, v_0). Suppose $(u_0, v_0) \in E(C_0)$ (otherwise choosing an edge in $E(C_0)$ replace for (u_0, v_0)). Since there exists at most one faulty element outside BH_{n-1}^0, by Lemma 2.9(3), (u_0, v_0) is contained in a fault-free cycle C, say $C = \langle v_0, u_0, v_1, u_1, v_2, u_2, v_3, u_3, v_0 \rangle$, of length 8 in BH_n such that $E(C) \cap E(BH_{n-1}^i) = \{(u_i, v_i)\}$ for each $i \in \{0,1,2,3\}$. Since $f_t \leq 1$ for all $t \in \{1,2,3\}$, note that $1 \leq n-1$ and $1 \leq 2n-3$ for $n \geq 3$, by Lemma 2.12 or Lemma 2.11, there exists a fault-free cycle C_t of length $2^{2(n-1)} - 2f_v^t$ passing through (u_t, v_t). Let $P_i = C_i - (u_i, v_i)$ for $i \in \{0,1,2,3\}$, then $\langle u_0, P_0, v_0, u_3, P_3, v_3, u_2, P_2, v_2, u_1, P_1, v_1, u_0 \rangle$ is a desired fault-free cycle of length $2^{2n} - 2f_v$ in BH_n.

Subcase 2.2. $f_v^0 = n-1$.

It implies $f_v^j = 0$ and $f_e^j \leq 1$ for each $j \in \{1,2,3\}$. We regard one faulty vertex in BH_{n-1}^0 as fault-free temporarily. For the situation where $f_0 - 1 \leq 2n-4$ and $f_v' = f_v^0 - 1 = f_v - 1 = n-2$, by the inductive hypothesis, there exists a cycle C_0 in BH_{n-1}^0 of length $2^{2(n-1)} - 2f_v'$, where $f_v' = f_v^0 - 1$. Note that C_0 contains at most one faulty vertex.

If C_0 does not contain any faulty vertex, choose an edge, say $(u_0, v_0) \in E(C_0)$ (assume u_0 is white, v_0 is black). By Lemma 2.9(3) and $f - f_0 \leq 1$, (u_0, v_0) is contained in a fault-free cycle C, say $C = \langle v_0, u_0, v_1, u_1, v_2, u_2, v_3, u_3, v_0 \rangle$, of length 8 in BH_n such that $E(C) \cap E(BH_{n-1}^i) = \{(u_i, v_i)\}$ for each $i \in \{0,1,2,3\}$. By using the method similar to that of Subcase 1.2.2, we can find a desired cycle of length $2^{2n} - 2f_v$ in BH_n. The details are omitted.

Now we assume that C_0 contains one faulty vertex u_0 and u_0 is black (if u_0 is white, the discussions are similar). Let a_0 and b_0 be two vertices adjacent to u_0 in C_0. Let $c_0 \in C_0$ be a neighbor of b_0. Recall that each

vertex has two extra neighbors and $f_c \leq 1$, we can choose the extra neighbors $c_1 \in V(BH^1_{n-1})$ and $a_3 \in V(BH^3_{n-1})$ of a_0 and c_0, respectively, such that both (a_0, c_1) and (c_0, a_3) are fault-free. The desired cycle of BH_n can be constructed by the similar discussions as Subcase 1.2.2, the details are omitted.

Case 3. There exists one $i \in \{0, 1, 2, 3\}$ such that $f_i = 2n - 2$.

Without loss of generality, let $i = 0$, i.e. $f_0 = f = 2n - 2$. It implies that $f_j = 0$ for all $j \in \{1, 2, 3\}$ and $f_c = 0$.

Since $f_v \geq 1$, we can choose a faulty vertex, say w (assume w is black, the discussion is similar for w being white). Note that $f_v \leq n - 1$, $f_e = f - f_v \geq 2n - 2 - (n - 1) = n - 1 \geq 1$ for $n \geq 3$, let (u_0, v_0) (assume v_0 is black) be a faulty edge. Image w and (u_0, v_0) as fault-free temporarily. Since $f_0 - 2 \leq 2n - 4$ and $f'_v = f^0_v - 1 \leq n - 1 - 1 = n - 2$, by the inductive hypothesis, there exists a cycle C_0 in BH^0_{n-1} of length $2^{2(n-1)} - 2f'_v$, where $f'_v = f^0_v - 1 = f_v - 1$. Note that C_0 contains at most one faulty vertex w and one faulty edge (u_0, v_0). We consider the following four subcases.

Subcase 3.1. $w \notin C_0$ and $(u_0, v_0) \notin E(C_0)$.

Choose any edge, say $(a_0, b_0) \in E(C_0)$ (assume a_0 is white, b_0 is black). By Lemma 2.9(1) and $f_c = 0$, (a_0, b_0) is contained in a fault-free cycle C, say $C = \langle b_0, a_0, b_1, a_1, b_2, a_2, b_3, a_3, a_0\rangle$, of length 8 in BH_n such that $E(C) \cap E(BH^i_{n-1}) = \{(a_i, b_i)\}$ for each $i \in \{0, 1, 2, 3\}$. Since for any $i \in \{1, 2\}$, $f_i = 0$, by Lemma 2.5, there exists a fault-free path P_i of length $2^{2(n-1)} - 1$ between a_i and b_i in BH^i_{n-1} for $i = 1, 2$. Since $f_3 = 0$, by Lemma 2.6, there exists a fault-free path P_3 of length $2^{2(n-1)} - 3$ between a_3 and b_3 in BH^i_{n-1}. Let $P_0 = C_0 - (a_0, b_0)$. A desired fault-free cycle of $2^{2n} - 2f_v$ in BH_n can be constructed as $\langle a_0, b_1, P_1, a_1, b_2, P_2, a_2, b_3, P_3, a_3, b_0, P_0, a_0\rangle$.

Subcase 3.2. $w \notin C_0$ and $(u_0, v_0) \in E(C_0)$.

By Lemma 2.9(1) and $f_c = 0$, (u_0, v_0) is contained in a cycle C, say $C = \langle v_0, u_0, v_1, u_1, v_2,$
$u_2, v_3, u_3, v_0\rangle$, of length 8 in BH_n such that $E(C) \cap E(BH^i_{n-1}) = \{(u_i, v_i)\}$ and (u_i, v_{i+1}) is fault-free for each $i \in \{0, 1, 2, 3\}$. A desired cycle can be constructed by the similar discussion as Subcase 3.1, the details are omitted.

Subcase 3.3. $w \in C_0$ and $(u_0, v_0) \notin E(C_0)$.

Let a_0 and b_0 be two vertices adjacent to w in C_0. Since each vertex has two extra neighbors, let $a_1 \in V(BH^1_{n-1})$, $b_1 \in V(BH^1_{n-1})$ be an extra neighbor of a_0 and b_0, respectively, such that $a_1 \neq b_1$. Choose two distinct white vertices, say c_i, d_i, respectively, in BH^i_{n-1} for $i = 1, 2$. Let one of the extra neighbors of c_i, d_i be e_{i+1}, g_{i+1} in BH^{i+1}_{n-1}, respectively, for $i = 1, 2$. See Figure 4. By Lemma 2.8, there exist two vertex-disjoint paths $P_1[a_1, c_1]$ and $R_1[b_1, d_1]$ (resp. $P_2[c_2, e_2]$ and $R_2[d_2, g_2]$) in BH^1_{n-1} (resp. BH^2_{n-1}) such that $V(P_1[a_1, c_1]) \cup V(R_1[b_1, d_1]) = V(BH^1_{n-1})$ (resp. $V(P_2[c_2, e_2]) \cup V(R_2[d_2, g_2]) = V(BH^2_{n-1})$). By Lemma 2.7, there exists a



Figure 4: Illustration of 3.3;



Figure 5: Illustration of 3.4.2

path P_3 in $V(BH^3_{n-1})$ of length $2^{2(n-1)}-2$ between e_3 and g_3. Since $f_c = 0$, the crossing edges (c_1, e_2), (d_1, g_2), (c_2, e_3), (d_2, g_3) are fault-free. Let $P_0 = C_0 - \{w\}$. Thus, a fault-free cycle of length $2^{2n} - 2f_v^0 = 2^{2n} - 2f_v$ in BH_n is $\langle a_0, a_1, P_1[a_1, c_1], c_1, e_2, P_2[e_2, c_2], c_2, e_3, P_3, g_3, d_2, R_2[d_2, g_2], g_2, d_1, R_1[d_1, b_1], b_1, b_0, P_0, a_0\rangle$.

Subcase 3.4. $w \in C_0$ and $(u_0, v_0) \in E(C_0)$.

Let a_0 and b_0 be two vertices adjacent to w in C_0. If $v_0 = w$, then $u_0 = a_0$ or $u_0 = b_0$. Without loss of generality, assume that $u_0 = b_0$. The discussion is the same as that in Subcase 3.3.

Now we assume $v_0 \neq w$ in the following. Notice that u_0 may equal to a_0 or b_0.

Subcase 3.4.1. $u_0 = b_0$ (If $u_0 = a_0$, the discussion is similar).

Assume that $C_0 = \langle a_0, w, u_0, v_0, P_0[v_0, a_0], a_0\rangle$. For $i \in \{1, 2, 3\}$, let $u_i \in V(BH^i_{n-1})$ be a white vertex such that $(u_3, v_0) \in E(BH_n)$. For $j \in \{1, 2, 3\}$, let $v_j \in V(BH^j_{n-1})$ be a black vertex such that $(a_0, v_1), (u_1, v_2), (u_2, v_3) \in E(BH_n)$. Since $f_i = 0$ for $i \in \{1, 2, 3\}$, by Lemma 2.5, there exists a $u_i - v_i$ path P_i of length $2^{2(n-1)}-1$ in BH^i_{n-1}. Thus, $\langle a_0, v_1, P_1, u_1, v_2, P_2, u_2, v_3, P_3, u_3, v_0, P_0[v_0, a_0], a_0\rangle$ is a desired fault-free cycle of length $2^{2n} - 2f_v$ in BH_n.

Subcase 3.4.2. $u_0 \neq \{a_0, b_0\}$.

Let $c_1 \in V(BH^1_{n-1})$ (resp. $u_3 \in V(BH^3_{n-1})$) be an extra neighbor of a_0 (resp. v_0). Let $c_0 \in C_0$ be a neighbor of b_0 such that $c_0 \neq v_0$ (otherwise let $c_0 \in C_0$ be a neighbor of a_0, the discussion is similar). Thus C_0 is the type as $C_0 = \langle a_0, w, b_0, c_0, P_0[c_0, u_0], u_0, v_0, P_0[v_0, a_0], a_0\rangle$ (to see Figure 5). Since each vertex has two extra neighbors, let $v_1 \in V(BH^1_{n-1})$ (resp. $a_3 \in V(BH^3_{n-1})$) be an extra neighbor of u_0 (resp. c_0) such that $v_1 \neq c_1$ (resp. $a_3 \neq u_3$). For $i \in \{1, 2\}$, let $u_i, a_i \in V(BH^i_{n-1})$ be two dis-

tinct white vertices. For $j \in \{2,3\}$, let $v_j, c_j \in V(BH^j_{n-1})$ be two distinct black vertices such that $(u_{j-1}, v_j) \in E(BH_n)$ and $(a_{j-1}, c_j) \in E(BH_n)$. Since $f_i = 0$ for $i \in \{1,2,3\}$, by Lemma 2.8, there exist two vertex-disjoint paths $P_i[v_i, u_i]$ and $R_i[c_i, a_i]$ in BH^i_{n-1} such that $V(P_i[v_i, u_i]) \cup V(R_i[c_i, a_i]) = V(BH^i_{n-1})$. Thus, a fault-free cycle of length $2^{2n} - 2f_v$ in BH_n is $\langle a_0, c_1, R_1[c_1, a_1], a_1, c_2, R_2[c_2, a_2], a_2, c_3, R_3[c_3, a_3], a_3, c_0, P_0[c_0, u_0], u_0, v_1, P_1[v_1, u_1], u_1, v_2, P_2[v_2, u_2], u_2, v_3, P_3[v_3, u_3], u_3, v_0, P_0[v_0, a_0], a_0\rangle$.

By the above cases, the proof is complete. □

4 Conclusion

In this paper, we obtained that BH_n with $|F_v| + |F_e| \leq 2n-2$ and $|F_v| \leq n-1$ has a fault-free cycle of length $2^{2n} - 2|F_v|$. If $|F_v| = 0$, $|F_e| \leq 2n-2$, a fault-free cycle of length $2^{2n} - 2|F_v| = 2^{2n}$ is a Hamiltonian cycle. Since BH_n is a bipartite graph with two partite sets of equal size, the cycle is the longest in the worst-case. Furthermore, since the edge connectivity of BH_n is $2n$, the BH_n cannot tolerate $2n-1$ faulty edges. Hence, our result is optimal in terms of faulty edges.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No.11371052, No.11571035, No.11271012 and No. 61572005).

References

[1] B. Alspach, D. Hare, Edge-pancyclic block-intersection graphs, Discrete Math. 97 (1991) 17–24.

[2] J.A. Bondy, Pancyclic graphs, J. Combin. Theory 11 (1971) 80–84.

[3] D. Cheng, R.-X. Hao, Various cycles embedding in faulty balanced hypercubes, Inform. Sci. 297 (2015) 140–153.

[4] D. Cheng, R.-X. Hao, Y.-Q. Feng, Vertex-fault-tolerant cycles embedding in balanced hypercubes, Inform. Sci. 288 (2014) 449–461.

[5] D. Cheng, R.-X. Hao, Y.-Q. Feng, Two node-disjoint paths in balanced hypercubes, Appl. Math. Comput. 242 (2014) 127–142.

[6] Z.-Z. Du, J.-M. Xu, A note on cycle embedding in hypercubes with faulty vertices, Inform. Proces. Lett. 111 (2011) 557–560.

[7] J.-S. Fu, Longest fault-free paths in hypercubes with vertex faults, Inform. Sci. 176 (2006) 759–771.

[8] R.-X. Hao, R. Zhang, Y.-Q. Feng, J.-X. Zhou, Hamiltonian cycle embedding for fault tolerance in balanced hypercubes, Appl. Math. Comput. 244 (2014) 447–456.

[9] S.-Y. Hsieh, Fault-tolerant cycle embedding in the hypercube with more both faulty vertices and faulty edges, Parall. Comput. 32 (2006) 84–91.

[10] K. Huang, J. Wu, Balanced hypercubes, in: Proc. of the 1992, Int. Conf. Parallel Process. 3 (1992) 153–159.

[11] K. Huang, J. Wu, Fault-tolerant resource placement in balanced hypercubes, Inform. Sci. 99 (1997) 159–172.

[12] K. Huang, J. Wu, Area efficient layout of balanced hypercubes, Int. J. High Speed Electron. Syst. 6 (1995) 631–646.

[13] H. Lü, X. Li, H. Zhang, Matching preclusion for balanced hypercubes, Theoret. Comput. Sci. 465 (2012) 10–20.

[14] H. Lü, H. Zhang, Hyper-Hamiltonian laceability of balanced hypercubes, J. Supercomput. 68 (1) (2014) 302–314.

[15] C.-H. Tsai, Fault-tolerant cycles embedded in hypercubes with mixed link and node failures, Appl. Math. Lett. 21 (2008) 855–860.

[16] J. Wu, K. Huang, The balanced hypercubes: a cube-based system for fault-tolerant applications, IEEE Trans. Comput. 46 (4) (1997) 484–490.

[17] J.-M. Xu, M. Ma, Survey on path and cycle embedding in some networks, Front. Math. China 4 (2009) 217–252.

[18] M. Xu, X.-D. Hu, J.-M. Xu, Edge-pancyclicity and Hamiltonian laceability of the balanced hypercubes, Appl. Math. Comput. 189 (2007) 1393–1401.

[19] M.-C. Yang, Bipanconnectivity of balanced hypercubes, Comput. Math. Appl. 60 (2010) 1859–1867.

[20] M.-C. Yang, Super connectivity of balanced hypercubes, Appl. Math. Comput. 219 (2012) 970–975.

[21] M.-C. Yang, Conditional diagnosability of balanced hypercubes under the PMC model, Inform. Sci. 222 (2013) 754–760.

[22] M.-C. Yang, J.J.M. Tan, L.H. Hsu, Hamiltonian circuit and linear array embeddings in faulty k-ary n-cubes, J. Parall. Distrib. Comput. 67 (4) (2007) 362–368.

[23] M.-C. Yang, M.-H. Yang, Reliability analysis of balanced hypercubes, IEEE COMCOMAP, (2012) 376–379.

[24] W.-H. Yang, H.Z. Li, W.H. He, Edge-fault-tolerant bipancyclicity of Cayley graphs generated by transposition-generating trees, Inter. J. Comput. Math. 92(2015) 1345–1352.

[25] J.-X. Zhou, Z.-L. Wu, S.-C. Yang, K.-W. Yuan, Symmetric property and reliability of balanced hypercube, IEEE Trans. Comput. 64 (2015) 876–881.

[26] Q. Zhou, D. Chen, H. Lü, Fault-tolerant Hamiltonian laceability of balanced hypercubes, Inform. Sci. 300 (2015) 20–27.